


Supplementary Figure 1. Putative binding-site consensus logos for different regulons. The logos represent the consensus sequence generated with for the putative DNA binding-sites for each regulon. The sequence underneath the diagram represents previously reported consensus binding site for each family. Logos were generated by WebLogo algorithm using the entire set of binding site sequences identified for each *E. faecalis* transcription family.



References

DnaA

Hansen FG, Christensen BB, Nielsen CB, Atlung T. Insights into the quality of DnaA boxes and their cooperativity. *J Mol Biol.* 2006 Jan 6;355(1):85-95.

Fis

Stella S, Cascio D, Johnson RC. The shape of the DNA minor groove directs binding by the DNA-bending protein Fis. *Genes Dev.* 2010 Apr 15;24(8):814-26.

LexA, ArgR and Pho

Li H, Rhodius V, Gross C, Siggia ED. Identification of the binding sites of regulatory proteins in bacterial genomes. *Proc Natl Acad Sci U S A.* 2002 Sep 3;99(18):11772-7.

DeoR

Zeng X, Saxild HH. Identification and characterization of a DeoR-specific operator sequence essential for induction of *dra-nupC-pdp* operon expression in *Bacillus subtilis*. *J Bacteriol.* 1999 Mar;181(6):1719-27.

Mall

Lloyd GS, Godfrey RE, Busby SJ. Targets for the Mall repressor at the divergent *Escherichia coli* K-12 *malX-mall* promoters. *FEMS Microbiol Lett.* 2010 Apr;305(1):28-34.

GalR-S, PurR, FruR and CRP-FNR

Gelfand MS, Novichkov PS, Novichkova ES, Mironov AA. Comparative analysis of regulatory patterns in bacterial genomes. *Brief Bioinform.* 2000 Nov;1(4):357-71.

Fur

Fuangthong M, Helmann JD. Recognition of DNA by three ferric uptake regulator (Fur) homologs in *Bacillus subtilis*. *J Bacteriol.* 2003 Nov;185(21):6348-57.

CopY

Portmann R, Magnani D, Stoyanov JV, Schmechel A, Multhaup G, Solioz M.


Interaction kinetics of the copper-responsive CopY repressor with the *cop* promoter of *Enterococcus hirae*. *J Biol Inorg Chem.* 2004 Jun;9(4):396-402.

LysR


Schell, M. 1993. Molecular biology of the LysR family of transcriptional regulators. *Annu. Rev. Microbiol.* 47:597–626.

Grob P, Kahn D, Guiney DG. Mutational characterization of promoter regions recognized by the *Salmonella dublin* virulence plasmid regulatory protein SpvR. *J Bacteriol.* 1997 Sep;179(17):5398-406.

Supplementary Figure 2. Distribution of the position of the binding sites within the promoter region in *E. faecalis*. Bars indicate the total number of sequences contained within the promoter regions, grouped by their distance to the translational start site. Values are expressed as the percent of binding sites in each position range related to the total of binding sites within the network. (n = 629).



Supplementary Figure 3. Time course of intracellular copper content in *E. faecalis* wild type and *copΔ* strains grown in basal medium. The basal concentration of copper quantified was 10 μM . No significant differences between each column were detected. ANOVA test $p < 0.05$. Control time indicated exactly the moment was the bacteria were added in the media.



Supplementary Figure 4. Growth curve of wild type and *copΔ* strains exposed to different copper concentrations. White and black figures represent the wild type and *copΔ* strains, respectively. Growing medium: Basal N (circles), 0.5 mM CuSO₄ (squares) and 1 mM CuSO₄ (triangles). Each value corresponds to the average of 3 independent biological replicates (error bars indicate standard deviation).

